Advancing the Science of Air Monitoring:

Research Priorities and New Approaches for Enhancing Monitoring Data

Tim Watkins
National Exposure Research Laboratory
Office of Research and Development

National Ambient Air Monitoring Conference November 9, 2006

Presentation Overview

- Monitoring Objectives and Research Priorities
- Monitoring Research in EPA's Office of Research and Development (ORD)
- Approaches for Enhancing Ambient Monitoring



Monitoring for Multiple Objectives

Monitoring Purpose	Measurements Needed		
Compliance with NAAQS	FRM / FEM		
Public Information/Reporting	Continuous / real time measurements		
 Air Quality Management Activities Development of Emission Reduction Strategies Evaluation of Emission Reduction Strategies Air Quality Trends 	Continuous / semi-continuous sampling		
	PM Speciation		
	Precursor species		
	Air Toxics		
Research/Scientific Studies - Health Assessments - Ecological Assessment - Source-Receptor Relationships - Model Development and Evaluation	Continuous / semi-continuous sampling		
	PM Speciation		
	Precursor species and co-pollutants		
	Air Toxics		
	Deposition		



Monitoring Priorities to Support EPA Programs

- NAAQS Development
 - PM Size Fractions
 - Coarse and Ultrafine
 - PM Speciation / Components
- NAAQS Implementation
 - FRM / FFM
 - Emission Reduction Strategies Development
 - Accountability
- Public Reporting
 - AirNOW and AQI
- National Ambient Air Monitoring Strategy (Dec 05 Draft)
 - Continuous Measurements
 - Multipollutant Measurements (NCORE Sites)
 - Increased integration with Science Objectives
 - Integration of Air Toxics into monitoring networks
 - Ecological Assessment



Monitoring Priorities from External Recommendations: National Research Council Reports

- Research Priorities for Airborne Particulate Matter IV (2004)
 - Challenge Enhancing Air Quality Monitoring for Research
 - Move from a focus on compliance with NAAQS toward multiple monitoring purposes
 - Continuous / semi-continuous measurements
 - PM species
 - The report specifically mentions the need to measure ultrafine PM, soluble metals, and organic species.
- Air Quality Management in the US (2004)
 - Recommendation Enhance Air Pollution Monitoring
 - Expanded to other important objectives beyond compliance
 - Use of semi-continuous methods
 - Develop more reliable methods and analytical procedures for chemical composition of PM



Monitoring Priorities from External Recommendations: NARSTO PM Assessment (2004)

- Improve integration of atmospheric and health sciences to understand relationships between PM and public health impacts.
 - "Health impacts based on epidemiological studies can only be derived for PM characteristics for which ambient measurements are available."
- Improve the understanding of carbonaceous aerosols
 - Chemical speciation
 - Spatial and temporal resolution
 - Composition
- Develop methods to identify important markers or tracer species to relate ambient concentrations to sources.
- Replace integrated measurements with continuous, real-time measurements for PM mass and composition, where feasible and as technology evolves.





ORD Monitoring Research

- Methods Development and Evaluation
- Source Apportionment Applications
- Exposure Assessment Applications
- Health Effects Applications



ORD's Monitoring Research Program: Methods Development and Evaluation

- Sampling Methods
 - Coarse particles
 - Semi-continuous PM species
 - Air Toxics
 - Acrolein and 1,3 butadiene
 - Mercury dry deposition
- Analytical Methods
 - Inorganic and Organic Source marker compounds
- Research Grants
 - Carbonaceous PM
 - Source Apportionment
 - Continuous PM
- PM Supersites




ORD's Monitoring Research Program: Source Apportionment and Exposure Applications

- Detroit Exposure and Aerosol Research Study (DEARS)
 - PM Components and Size Fractions
 - Air Toxics
- Steubenville Source Apportionment Study
 - Integrated and Continuous PM
 - Continuous Criteria Gases
 - Deposition
 - Ambient Mercury Speciation
- Coarse PM Methods Evaluation
 - FRM / FEM
 - Continuous Methods
 - Saturation Samplers
- Near Roadway Exposures
 - PM Composition and Size Fractions
 - Mobile Source Air Toxics



Preliminary DEARS Results

(Concentration Ratios – Outdoor Residential to Community Central Site)




A STATE OF THE STA

Regional

Traffic

Diesel





Industrial

Steubenville PMF Apportionment Results 2003 & 2004

Analyte	Factor 1	Factor 2	Factor 3	Factor 4	Factor 5	Factor 6
Oil/	Incineration		Crustal		Iron/Steel	Coal
Mg	*	105.12	568.88	29.24	191.09	*
Al	76.43	*	365.66	38.50	56.26	38.94
P	*	65.72	*	*	*	*
S	*	*	534.38	*	*	11832.0
C1	2167.7	*	*	712.45	263.86	709.84
V	1.11	*	*	*	2.87	*
Cr	0.53	*	*	*	2.44	*
Mn	*	16.36	37.42	*	53.67	*
Fe	118.82	28.71	25.63	*	343.61	36.17
Ni	3.54	0.55	*	*	*	*
Cu	14.17	2.26	*	8.68	*	20.25
Zn	45.88	3.85	*	14.83	3.99	16.15
As	0.58	0.03	0.07	0.31	0.04	0.76
Se	0.22	0.02	*	1.39	*	2.45
Rb	0.21	0.27	0.14	0.09	*	0.25
Sr	2.78	1.57	5.77	*	0.56	1.25
Mo	*	*	*	3.84	*	*
Cd	0.23	*	*	0.26	0.10	0.37
La	0.11	*	0.64	0.01	*	*
Ce	*	*	1.26	*	0.04	*
Hg	*	0.004	*	*	0.012	0.142
Pb	5.92	0.18	0.22	1.34	1.03	4.57
NO_3	6143.4	104.33	1010.7	*	*	6515.2
% Hg Explained	*	2	*	*	5	74

^{* =} Not Significant at 95% confidence interval

Source: Keeler et al, Environment Science and Technology, in press



ORD's Monitoring Research Program: Health Effects Applications


- Epidemiological
 - Detroit Children's Health Study
 - · Childhood asthma and mobile sources
 - Multi-city
 - Compositional differences in Air Quality
 - Chronic exposures to PM
- Toxicological
 - Source-specific effects
 - Coarse particles
- PM Research Centers
 - Harvard University
 - New York University
 - University of Washington
 - Rochester University
 - UCLA






Satellite Data

- Emerging source of air quality data
- Aerosol optical depth (AOD) used to estimate ground level concentrations
- Spatial and Temporal Gaps
 - Cloud cover
 - Reflective surfaces










12 Sept. 2002-A close-up of Houston shows many of the hourly PM2.5 monitors recorded 24 averages in excess of 40.5 ug/m3, (AQI>100). High AOD extends into a large portion of TX.





"Data Fusion"


- Combining monitoring data with other sources of air quality data (e.g., modeled output, satellite) to generate air quality surfaces
 - Capitalize on the strengths of monitoring data ("true" measure) and modeling data (spatial and temporal coverage)
 - Minimize weaknesses of each data source




Data Fusion Example – PM Concentrations in NE US (Feb 14, 2001)







Data fusion results:
Spatially and temporally resolved surface enhanced with ground truth data from monitors





Summary

- Enhanced monitoring for multiple objectives
- Monitoring Priorities include
 - Continuous methods
 - PM species and size fractions
 - Air toxics
 - Increased integration with science objectives
- ORD Monitoring Research
 - Methods Development and Evaluation
 - Source Apportionment, Exposure, and Health Applications
- Approaches are emerging to enhance, not replace, ambient monitoring
 - Satellites
 - Data Fusion



